

Measuring the trilinear neutral Higgs boson couplings in the MSSM at e^+e^- colliders

Charanjit K. Khosa*

Centre for High Energy Physics, Indian Institute of Science, Bangalore 560 012, India

E-mail: khosacharanjit@gmail.com

P. N. Pandita

Centre for High Energy Physics, Indian Institute of Science, Bangalore 560 012, India

E-mail: pandita@associates.iucaa.in

We consider the measurement of the trilinear couplings of the neutral Higgs bosons (H^0, h^0) in the minimal supersymmetric standard model (MSSM) at a high energy e^+e^- linear collider in the light of the discovery of a Higgs boson at the CERN Large Hadron Collider (LHC). We identify the state observed at the LHC with the lightest CP-even Higgs boson of the MSSM. We implement this constraint, as well as all the other relevant experimental constraints, on the parameter space of the MSSM in order to study the feasibility of measuring the trilinear couplings of the neutral Higgs bosons. For the measurement of trilinear couplings, we consider the multiple Higgs production processes. We delineate the regions of MSSM parameter space where the trilinear couplings of the neutral Higgs bosons could be measured at a high energy electron-positron collider.

*38th International Conference on High Energy Physics
3-10 August 2016
Chicago, USA*

*Speaker.

1. Introduction

The ATLAS and CMS experiments at the CERN Large Hadron Collider (LHC) have discovered a particle with mass about 125 GeV [1] whose properties are consistent with the Higgs boson of the standard model (SM). To confirm that this particle is the Higgs boson of the SM, one must measure its couplings to other particles as well as to itself. In the SM, the potential of the Higgs doublet ϕ which breaks the $SU(2) \times U(1)$ gauge symmetry is written as

$$V_{SM}^h = \lambda \left(\phi^2 - \frac{v^2}{2} \right)^2 = \frac{m_h^2}{2} h^2 + \lambda_{hhh}^{SM} \frac{h^3}{3!} + \lambda_{hhhh}^{SM} \frac{h^4}{4!} \quad ; \quad \phi = \frac{1}{\sqrt{2}} (0 \ v + h)^T, \quad (1.1)$$

where h is the physical Higgs boson, and $\lambda_{hhh}^{SM} = 3m_h^2/m_Z^2 = 5.6454$ (in units of $(\sqrt{2}G_F)^{1/2}m_Z^2 = 33.77$ GeV) and $\lambda_{hhhh}^{SM} = 3m_h^2/m_Z^4 = 0.00068$ GeV⁻² (in units of $(\sqrt{2}G_F)m_Z^4 = 1140.52$ GeV²) are the triple and quartic Higgs couplings, respectively. One must, therefore, measure these couplings in order to confirm the idea of the spontaneous breaking of the underlying gauge symmetry. Any deviation from the SM prediction for these couplings will signal the existence of new physics beyond SM. Supersymmetry (SUSY)[2] is the leading candidate for the new physics which stabilizes the Higgs mass against large radiative corrections. The supersymmetric version of the SM, known as minimal supersymmetric standard model (MSSM)[2] not only solves the naturalness problem of the SM, but also leads to gauge coupling unification at a large scale ($M_{GUT} = 2 \times 10^{16}$ GeV). Furthermore, the lightest supersymmetric particle, usually the lightest neutralino, could be a possible dark matter candidate in R -parity conserving models. In the MSSM, there are two Higgs doublets (H_2, H_1) with opposite hypercharge, which break the SM gauge symmetry, and there are five physical Higgs bosons (h^0, H^0, A^0, H^\pm). In this paper we address the question of the measurement of some of the trilinear couplings of the neutral Higgs bosons h^0, H^0 at a high energy electron-positron (e^+e^-) collider.

2. MSSM Higgs sector and trilinear couplings of h^0, H^0

In this section we shall consider the trilinear couplings of the CP-even Higgs bosons (h^0, H^0). At the tree level the masses of all the Higgs bosons and their couplings can be written in terms of the mass of the CP-odd Higgs boson m_{A^0} and $\tan \beta = \langle H_2^0 \rangle / \langle H_1^0 \rangle$. For the phenomenological survey, the Higgs sector of the MSSM can be divided into two regions known as non-decoupling regime and decoupling regime. In the first case, $m_{A^0} \leq 130$ GeV, and heavy Higgs (H^0) is considered as SM-like (observed) state. On the other hand, decoupling regime is represented by $m_{A^0} > 300$ GeV, and the light Higgs boson (h^0) is identified with the observed 125 GeV Higgs boson. We shall here focus on the decoupling scenario. The mass matrix for the CP-even Higgs bosons of the MSSM can be written as

$$\mathcal{M}^2 = \begin{bmatrix} m_{A^0}^2 \sin^2 \beta + m_Z^2 \cos^2 \beta & -(m_Z^2 + m_{A^0}^2) \sin \beta \cos \beta \\ -(m_Z^2 + m_{A^0}^2) \sin \beta \cos \beta & m_{A^0}^2 \cos^2 \beta + m_Z^2 \sin^2 \beta \end{bmatrix} + \frac{3g^2}{16\pi^2 m_W^2} \begin{bmatrix} \Delta_{11} & \Delta_{12} \\ \Delta_{12} & \Delta_{22} \end{bmatrix}, \quad (2.1)$$

where the radiative corrections Δ_{ij} are sensitive, besides other parameters, to the top-stop masses. Other parameters that enter the calculations are the Higgs(ino) parameter μ , and $\tan \beta$. We have used CalcHEP [3] for the numerical calculations of the Higgs spectrum. Furthermore, we have adjusted the trilinear parameter A_t so as to fix the mass of the lightest Higgs (m_{h^0}) in the range 122 – 128 GeV. In Fig. 1 (left panel) we show the variation of the CP-even heavy Higgs mass (m_{H^0}) as a function of the parameter μ for fixed values of the supersymmetry breaking scale (M_S) and

$\tan\beta$. We see from this Fig. that the heavy Higgs mass is weakly dependent on M_S for fixed value of $\tan\beta$. Now coming to the Higgs self couplings, assuming CP -conservation, we have six couplings amongst the neutral Higgs bosons of the MSSM. These are written as λ_{hhh} , λ_{Hhh} , λ_{hAA} , λ_{HAA} , λ_{HHH} and λ_{HHh} . As in the case of the masses of the Higgs bosons, these trilinear couplings also obtain significant radiative corrections [4]. We can, then, write these trilinear couplings generically as $\lambda = \lambda^0 + \Delta\lambda$ where λ^0 is the tree-level coupling and $\Delta\lambda$ are the radiative correction. In this study we shall consider only two of these trilinear couplings, namely (in units of $(\sqrt{2}G_F)^{1/2}m_Z^2$) $\lambda_{hhh}^0 = 3\cos 2\alpha \sin(\beta + \alpha)$ and $\lambda_{Hhh}^0 = 2\sin 2\alpha \sin(\beta + \alpha) - \cos 2\alpha \cos(\beta + \alpha)$, where α is the mixing angle in the CP -even Higgs sector. These couplings are sensitive to m_{A^0} values of upto 500 GeV. The dependence of the trilinear couplings on the parameter μ is shown in Fig.1.




Figure 1: The mass of heavier Higgs boson m_{H^0} as a function of μ parameter for fixed $\tan\beta$ and M_S (left panel); variation of radiatively corrected trilinear couplings in MSSM with μ parameter(central panel); Feynman diagrams for resonant production of hh , through $e^+e^- \rightarrow HZ, HA, \nu_e \bar{\nu}_e H$ (where $H \rightarrow hh$ in the final state) and through non-resonant WW fusion (right panel).

3. Measurement of trilinear couplings

We now discuss the multiple Higgs production processes which can be used to study the trilinear couplings involving H^0 and h^0 . We consider the production of heavy Higgs boson through Higgs-strahlung $e^+e^- \rightarrow ZH$, associated production with CP -odd Higgs boson $e^+e^- \rightarrow AH$, and WW fusion mechanism $e^+e^- \rightarrow \nu_e \bar{\nu}_e H$ (see Fig. 1 (right panel) for Feynman diagrams and [4] for cross-sections). Multiple light Higgs bosons (h^0) can be produced through heavy CP -even Higgs boson decay. Non- resonant hh pair could also be produced with Z or A [4]. Associated production of A with h , where A decays to hZ , contributes as the background to the multiple Higgs production processes. In Fig. 2, left panel, we show the cross-section for $e^+e^- \rightarrow ZH, AH, H\nu\bar{\nu}$ as a func-




Figure 2: Cross-section for $e^+e^- \rightarrow AH, ZH, Ah/2, H\nu\bar{\nu}$ as the function of m_{H^0} for $\sqrt{s} = 500$ GeV (left); the branching fractions for different decay channels of H for both the benchmark points(central and right).

tion of m_{H^0} for $\sqrt{s} = 500$ GeV. We choose two benchmark values for the parameters $(\mu, M_1, M_2) = (-230, 120, 240)$ and $(-500, 150, 300)$, respectively, so as to include SUSY particles in the final

state of the heavy Higgs boson decay. In Fig. 2, central and right panel, we show the branching fractions of heavy Higgs decay into different channels. Second benchmark values have larger $\text{BR}(H \rightarrow hh)$ because neutralino and chargino spectrum is heavy as compared to the first benchmark point. Consequently, $\text{BR}(H \rightarrow \chi^0 \chi^0, \chi^+ \chi^-)$ is suppressed. In Fig. 3, for the measurement of λ_{Hhh} coupling, we show the contours of $\sigma(H) \times \text{BR}(H \rightarrow hh)$ for $\sqrt{s} = 500$ GeV and $\sqrt{s} = 1.5$ TeV, respectively for the first benchmark point. The decay $H \rightarrow hh$ is kinematically forbidden for $m_{A^0} \approx m_{H^0} \leq 250$ GeV, and this branching ratio decreases as we move diagonally upward in the $(m_{A^0}, \tan\beta)$ plane. The lower left corner of $(m_{A^0}, \tan\beta)$ plane is the suitable region to measure λ_{Hhh} coupling. The non-resonant WW fusion to $hh\nu_e\bar{\nu}_e$ final state involves both λ_{hhh} and λ_{Hhh} couplings. Having an estimate of λ_{Hhh} , we use this process to measure λ_{hhh} coupling. We can see from the Fig. 3, right panel, that the $\sigma(hh\nu_e\bar{\nu}_e)$ is almost independent of m_{A^0} and $\tan\beta$.




Figure 3: The contours of constant $\sigma(H) \times \text{BR}(H \rightarrow hh)$ (in fb) (left and central panel); the contours of non-resonant $\sigma(ee \rightarrow hh\nu\bar{\nu})$ (in fb) via non-resonant WW fusion (right).

4. Conclusions

We have studied the trilinear couplings of the CP-even Higgs bosons (H^0, h^0) in the minimal supersymmetric standard model, and their measurement at a high energy e^+e^- collider. In doing so we have identified the resonance observed at the CERN LHC at 125 GeV with the lightest CP even Higgs boson of the MSSM. By estimating the production cross-section of the various processes involving multiple Higgs bosons, we have delineated the regions in the $(m_{A^0}, \tan\beta)$ space where trilinear couplings λ_{Hhh} and λ_{hhh} can be measured at an e^+e^- collider. For the coupling λ_{Hhh} , we have $\sigma(H) \times \text{BR}(H \rightarrow hh) \approx 0.005$ fb and 0.04 fb for $\sqrt{s} = 500$ GeV and $\sqrt{s} = 1.5$ TeV, respectively, and it decreases along the diagonal upward direction in the $(m_{A^0}, \tan\beta)$ plane. We estimate the coupling λ_{hhh} through non-resonant WW fusion process. Precise knowledge of neutralino and chargino masses is crucial in order to determine the trilinear couplings of H^0, h^0 of the MSSM.

References

- [1] G. Aad *et al.* [ATLAS and CMS Collaborations], *Combined Measurement of the Higgs Boson Mass in pp Collisions at $\sqrt{s} = 7$ and 8 TeV with the ATLAS and CMS Experiments*, *Phys. Rev. Lett.* **114** 191803 (2015).
- [2] H. P. Nilles, *Supersymmetry, Supergravity and Particle Physics*, *Phys. Rept.* **110** 1 (1984).
- [3] A. Belyaev, N. D. Christensen and A. Pukhov, *CalcHEP 3.4 for collider physics within and beyond the Standard Model*, *Comput. Phys. Commun.* **184** 1729 (2013).
- [4] P. Osland and P. N. Pandita, *Measuring the trilinear couplings of MSSM neutral Higgs bosons at high-energy e^+e^- colliders*, *Phys. Rev. D* **59** 055013 (1999) ; [hep-ph/9911295].
- [5] C. K. Khosa and P. N. Pandita, *Measuring the trilinear neutral Higgs boson couplings in the minimal supersymmetric standard model at e^+e^- colliders in the light of the discovery of a Higgs boson*, *Int. J. Mod. Phys. A* **31** 1650108 (2016), and references therein.